

Metrics Fluctuational Theory

Timur. F. Kamalov

Physics Department, Moscow State Opened University,
ul. P. Korchagina, 22, Moscow 107996, Russia

Tel./fax 7-095-2821444

E-mail: okamalov@chat.ru

ykamalov@rambler.ru

It is supposed the alternative to Quantum Mechanics Axiomatic. Fluctuational Theory save the Mathematics of Quantum Mechanic without change, naming this Mathematics as Method of Indirect Computation.

Fluctuational Theory is delete the axiomatic of Quantum Mechanics and replaces it by the assumption of Gravitational Noise. This assumption is connects the Method of Indirect Computation to the Classical Physics.

Physical fluctuations of classical gravitational fields are mathematically expressed through geometric fluctuations of metrics of Riemann Space.

Metrics Fluctuational Theory and Quantum Mechanic is describe the classical experiment of electrons interference by two different way.

Pacs 03.65*

Key words: Fluctuational Theory, Gravity Background, Gravity Noise, Polarization Variables, Entanglement State.

Introduction

Classical particles in stochastic classical gravitational fields called here the microobjects in random gravitational background. The latter is assumed to be the background of stochastic gravitational fields and waves distributed isotropically over the space-time in the average. The concept of probability interval in the Riemann space is introduced. The interference experiment with electrons is described here. Physical fluctuations of classical gravitational fields are mathematically expressed through geometric fluctuations of metrics of 4-dimensional Riemann space.

Main part

Quantum mechanics is the theory with axiomatic is based on basic results obtained in a number of experiment studies. These experiments were impossible to explain within the framework of classical physics, and they constituted the foundations of the new theory called quantum one.

Though this new theory did not investigate the nature of experiments laid in its foundation, as neither theory would investigate its own axioms, nevertheless, it enabled performing specific calculations of real physical experiments and was capable of correctly predicting the behavior of quantum micro-objects. Let us start considering these experiments from the viewpoint of classical physics, and further pass to their discussion with the help of quantum theory.

Let us agree to consider the physical gravitational field really existing, its approximate mathematical description to be expressed through geometrical fluctuations of spatial metrics. This spares us difficulties with energy-momentum conservation, as we consider the physical field to transfer energy - in contrast with the mathematical description of gravitational effects with Riemann geometry where the geometry is usually considered not to contain and not to transfer energy-momentum. In this, we shall confine ourselves to the linear approximation of metrics decomposition, which describes rather comprehensively all known experiments in verification of the General Relativity Theory.

Let us consider two classic particles in a field of random gravitational fields or waves. The General Theory of Relativity gives the length element in 4-dimensional Riemann space as

$$\Delta\ell^2 = g_{ik}\Delta x^i\Delta x^k,$$

the metric in the linear approach is

$$g_{ik} = \eta_{ik} + h_{ik},$$

being η_{ik} Minkowsky metric, constituting the unity diagonal matrix. Hereinafter, the indices μ, ν, γ acquire values 0, 1, 2, 3. Indices encountered twice imply summation thereupon. Let us select harmonic coordinates (the condition of harmonically coordinates mean selection of concomitant frame $\frac{\partial h_{\mu\nu}^m}{\partial x^m} = \frac{1}{2} \frac{\partial h_{\mu\nu}^m}{\partial x^\mu}$) and let us take into consideration that $h_{\mu\nu}$ satisfies the gravitational field equations

$$\square h_{mn} = -16\pi G S_{mn},$$

which follow from the General Theory of Relativity; here S_{mn} is energy-momentum tensor of gravitational field sources with d'Alembertian $\square$ and gravity constant G . Then, the solution shall acquire the form

$$h_{\mu\nu} = e_{\mu\nu} \exp(ik_\gamma x^\gamma) + e_{\mu\nu}^* \exp(ik_\gamma x^\gamma),$$

where the value $h_{\mu\nu}$ is called metric perturbation, $e_{\mu\nu}$ polarization, and k_γ is 4-dimensional wave vector. We shall assume that , which constitute metric perturbation $h_{\mu\nu}$, are distributed in space with an unknown distribution function $\rho = \rho(h_{\mu\nu})$.

Relative oscillations ℓ of two particles in classic gravitational fields are described in the General Theory of Relativity by deviation equations

$$\frac{D^2}{D\tau^2} \ell^i = R_{kmn}^i \ell^m \frac{dx^k}{d\tau} \frac{dx^n}{d\tau},$$

being R_{kmn}^i gravitational field Riemann's tensor.

Specifically, the deviation equations give the equations for two particles oscillations

$$\ddot{\ell}^1 + c^2 R_{010}^1 \ell^1 = 0, \quad \omega = c\sqrt{R_{010}^1}.$$

The solution of this equation has the form

$$\ell^1 = \ell_0 \exp(k_a x^a + i\omega t),$$

being $a = 1, 2, 3$. Each resultant gravitational field or wave with index j and Riemann's tensor $R_{kmn}^i(j)$ shall be corresponding to the value $\ell^i(j)$ with stochastically modulated phase $\Phi(j) = \omega(j)t$.

We assume to be dealing with a stochastic gravitational background, so we are to define the probability in 4-dimensional Riemann space. This definition shall satisfy the following requirements.

We shall call the stochastic curved space the 4-dimensional Riemann space with probabilities defined therein. With this, we shall require the following:

1. If the interval $\Delta\ell$ in the Riemann space equals zero, then the probability of finding the particle in this interval equals unity.
2. If the interval $\Delta\ell$ in the Riemann space equals infinity, then the probability of finding the particle in this interval equals zero.
3. The probability to find the particle in the intervals $\Delta\ell(x_2^i, x_1^i) + \Delta\ell(x_3^i, x_2^i) \geq \Delta\ell(x_3^i, x_1^i)$, equals $P_{21} + P_{32} \leq P$.

Hereinafter, we refer to the regions of 4-dimensional Riemann space, which are bount with each other with single type intervals, that is, only with ether time-like intervals or space-like intervals.

It should be noted that these requirements are met by intervals with the following type of distribution: interval $\Delta\ell$ is corresponding with the probability interval

$$\Delta P = \frac{1}{\sigma\sqrt{2\pi}} \exp\left(-\frac{\Delta\ell^2}{2\sigma^2}\right).$$

Similarly, for the probability interval of finding a particle with velocity u^i we have:

$$\Delta P = \frac{1}{\sigma\sqrt{2\pi}} \exp\left(-\frac{S}{S_0}\right),$$

or

$$\Delta P = \frac{1}{\sigma\sqrt{2\pi}} \exp\left(-\frac{W}{m\sigma^2}\right),$$

being W the energy of oscillation of the micro-particle in question acquired in the gravitational background; for a free particle with the RMS velocity deviation $(\Delta u^i)^2$ this energy equals action function S . Assuming the dispersion to be characteristic for the gravitational background and formally equating the dispersion with gravitational background energy, we obtain

$$\Delta P = -\frac{1}{\sigma\sqrt{2\pi}} \exp\left(-\frac{S}{S_0}\right).$$

Then, introducing notations $a^2 = \frac{1}{\sigma\sqrt{2\pi}}$, $S_0 = \frac{m\sigma^2}{2}$ and taking into account that the probability is equal to squared probability amplitude, the probability amplitude, characterizing the action acquired by any micro-particle of mass m in the gravitational background, will equal

$$\psi = a \exp\left(i\frac{S}{S_0}\right).$$

The probability amplitude is a vector in the complex space. Therefore, it can be decomposed in vector basis e^m , so that

$$\psi = e^m \psi_m.$$

The scalar product of two vectors A^m and B^n in the Riemann space is defined as $g_{mn}A^mB^n$, hence we can normalize this wave function as follows:

$$\int g_{mn}\psi_m\psi_n dx = 1.$$

If we pass to the 3-dimensional space, having added the classical Jacobi - Hamilton equation

$$\frac{\partial S}{\partial t} + \frac{1}{2m}(\nabla S)^2 + U = 0$$

and continuity equation for probability density $\psi^2 = a^2$,

$$\frac{\partial a^2}{\partial t} + \text{div}(a^2 \frac{\nabla S}{m}) = 0,$$

we shall get the Schroedinger equation for this wave function

$$i2S_0 \frac{\partial \psi}{\partial t} = -\frac{4S_0^2}{2m} \Delta \psi + U(x, y, z)\psi.$$

We consider the physical model with the gravitational noise [i.e. with the background of gravitational fields and waves]. This means that we assume existence of fluctuations in gravitational waves and fields expressed mathematically by metric fluctuations.

Let us consider now the experiment with interference of two electrons on two slits. The electron interference experiment is the one in which it is impossible to determine the electron trajectory. Any attempt to determine the electron trajectory fails due to any infinitesimal affecting of an electron with the purpose of determination of its trajectory would alter the interference pattern. This is the first aspect. On the other hand, interaction of classical and stochastic fields and waves in such experiments is usually neglected. Such interactions must exist in compliance with the existing provisions of classical physics, and in particular, of the General Relativity Theory. Moreover, this is experimentally confirmed by the pre-quantum classical physics, hence, they require verification of their effect onto quantum micro-objects.

Let us review some provisions of the General Relativity Theory. We consider the motion of electrons from the source S to the screen through slits 1 and 2.

$$\begin{aligned} \langle x | s \rangle_1 &= g_{\alpha\alpha}(1) \langle x | s \rangle_{\alpha\alpha} = \delta_{\alpha\alpha} \langle x | s \rangle_{\alpha\alpha} + h_{\alpha\alpha}(1) \langle x | s \rangle_{\alpha\alpha}, \\ \langle x | s \rangle_2 &= g_{\alpha\alpha}(2) \langle x | s \rangle_{\alpha\alpha} = \delta_{\alpha\alpha} \langle x | s \rangle_{\alpha\alpha} + h_{\alpha\alpha}(2) \langle x | s \rangle_{\alpha\alpha}, \end{aligned}$$

where $g_{\alpha\alpha}(1) \neq g_{\alpha\alpha}(2)$. Due to the propagation difference between the two trajectories in space and time, the interference pattern is generated. In the stochastic curved space one needs not to know the electron trajectory. The interference pattern emerges due to the difference in metrics $g_{\alpha\alpha}(1)$ and $g_{\alpha\alpha}(2)$

In this experiment we observe not interference of electrons, as usually it is supposed, and, from the point of view of the Metrics Fluctuations Theory, it

is interference of the metrics fluctuation which are moving through two hole. Here, electrons play the role of a medium(environment). We should add in this process resonant properties of environment(medium), i.e. electrons.

Conclusion

Presently, there is very little reliable information about random gravitational background. The latter is not investigated either theoretically or experimentally[1-9]. However, one can expect that the due account of it is capable of providing answers for a number of questions about the nature of microobjects.

Acknowledgments

I'm thanks to Doctor Yuriy Svirko from Tokyo University, Department of Applied Physics; to Professor Andrew Whitaker from Queen's Belfast University, Physics Department and to Professor Yuiry Rybakov from People's Friendship University of Russia, department of Theoretical Physics for help and assistant.

References

- [1] T. F. Kamalov, Quantum Mechanics with Hidden Variables..., Deponent VINITI, 1983, 092-145 (in Russian).
- [2] T. F. Kamalov, Bells Inequalities and Rieman's Space, "On the 25 years Bell's Inequalities", Moscow, 1990, p.137 (in Russian).
- [3] T. F. Kamalov, Nature of Quantum Statistics, Deponent VINITI, 1989, 4615-B89.
- [4] T. F. Kamalov, Background of random gravitation waves, Deponent VINITI, 1989, 4615-B89.
- [5] T. F. Kamalov, Quantum Computer and Hidden Variables, Fizicheskaia Misl Russii, 1, 2001 (in Russian).

- [6] T. F. Kamalov, Reformed Bell's Inequalities in 4-dimension Rieman's Space, 10-th UK Conference on the Foundations of Physics, 10-14 Sept., 2001, Belfast.
- [7] T. F. Kamalov, Bell's Inequalities in 4-dimension Rieman's Space, E-print arXiv, quant-ph/0109153, 29 Sept. 2001.
- [8] T. F. Kamalov, Hidden Variables and the Nature of Quantum Statistics, Journal of Russian Laser Reseach, Oct. 2001,v. 22, 5, p. 475.
- [9] T. F. Kamalov, Gravitational Noise with Polarizations Variables, the Nature of Entanglement States and Metric Fluctuational Theory of Quantum Mechanics, E-print arXiv, quant-ph/0205079, 14 May 2002.